

A NEW IMPROVED ERROR BOUND FOR LINEAR COMPLEMENTARITY PROBLEMS FOR B -MATRICES*

LEI GAO[†] AND CHAOQIAN LI[‡]

Abstract. A new error bound for the linear complementarity problem when the matrix involved is a B -matrix is presented, which improves the corresponding result in [C.Q. Li et al., A new error bound for linear complementarity problems for B -matrices. Electron. J. Linear Al., 31:476-484, 2016]. In addition some sufficient conditions such that the new bound is sharper than that in [M. García-Esnaola and J.M. Peña. Error bounds for linear complementarity problems for B -matrices. Appl. Math. Lett., 22:1071-1075, 2009] are provided.

Key words. Error bound, Linear complementarity problem, B -matrix

AMS subject classifications. 15A48, 65G50, 90C31, 90C33

1. Introduction. Given an $n \times n$ real matrix M and $q \in R^n$, the linear complementarity problem (LCP) is to find a vector $x \in R^n$ satisfying

$$(1.1) \quad x \geq 0, Mx + q \geq 0, (Mx + q)^T x = 0$$

or to show that no such vector x exists. We denote this problem (1.1) by $LCP(M, q)$. The $LCP(M, q)$ arises in many applications such as finding Nash equilibrium point of a bimatrix game, the network equilibrium problems, the contact problems and the free boundary problems for journal bearing etc, for details, see [1, 5, 18].

It is well-known that the $LCP(M, q)$ has a unique solution for any vector $q \in R^n$ if and only if M is a P -matrix [5]. Here a matrix M is called a P -matrix if all its principal minors are positive. For the $LCP(M, q)$, one of the interesting problems is to estimate

$$(1.2) \quad \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty,$$

which can be used to bound the error $\|x - x^*\|_\infty$ [4], that is,

$$\|x - x^*\|_\infty \leq \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \|r(x)\|_\infty,$$

*Received by the editors on Month x, 201x. Accepted for publication on Month y, 201y Handling Editor: .

[†]School of Mathematics and Information Science, Baoji University of Arts and Sciences, Baoji, Shannxi, 721007, P. R. China (gaolei@bjwlxy.edu.cn).

[‡]School of Mathematics and Statistics, Yunnan University, Kunming, Yunnan, 650091, P.R. China (lichaoqian@ynu.edu.cn).

where x^* is the solution of the $\text{LCP}(M, q)$, $r(x) = \min\{x, Mx + q\}$, $D = \text{diag}(d_i)$ with $0 \leq d_i \leq 1$ for each $i \in N$, $d = [d_1, d_2, \dots, d_n]^T \in [0, 1]^n$, and the min operator $r(x)$ denotes the componentwise minimum of two vectors.

When the matrix M for the $\text{LCP}(M, q)$ belongs to P -matrices or some subclass of P -matrices, various bounds for (1.2) were proposed, e.g., see [2, 3, 4, 7, 8, 9, 10, 11, 12, 13, 14] and references therein. Recently, García-Esnaola and Peña in [10] provided an upper bound for (1.2) when M is a B -matrix as a subclass of P -matrices. Here, a matrix $M = [m_{ij}] \in R^{n,n}$ is called a B -matrix [6] if for each $i \in N = \{1, 2, \dots, n\}$,

$$\sum_{k \in N} m_{ik} > 0, \text{ and } \frac{1}{n} \left(\sum_{k \in N} m_{ik} \right) > m_{ij} \text{ for any } j \in N \text{ and } j \neq i.$$

THEOREM 1.1. [10, Theorem 2.2] *Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form*

$$(1.3) \quad M = B^+ + C,$$

where

$$(1.4) \quad B^+ = [b_{ij}] = \begin{bmatrix} m_{11} - r_1^+ & \cdots & m_{1n} - r_1^+ \\ \vdots & & \vdots \\ m_{n1} - r_n^+ & \cdots & m_{nn} - r_n^+ \end{bmatrix}, \quad C = \begin{bmatrix} r_1^+ & \cdots & r_1^+ \\ \vdots & & \vdots \\ r_n^+ & \cdots & r_n^+ \end{bmatrix},$$

and $r_i^+ = \max\{0, m_{ij} | j \neq i\}$. Then

$$(1.5) \quad \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \leq \frac{n-1}{\min\{\beta, 1\}},$$

where $\beta = \min_{i \in N} \{\beta_i\}$ and $\beta_i = b_{ii} - \sum_{j \neq i} |b_{ij}|$.

It is not difficult to see that the bound (1.5) will be inaccurate when the matrix M has very small value of $\min_{i \in N} \{b_{ii} - \sum_{j \neq i} |b_{ij}|\}$, for details, see [15, 16]. To conquer this problem, Li et al., in [17] gave the following bound for (1.2) when M is a B -matrix, which improves those provided by Li and Li in [15, 16].

THEOREM 1.2. [17, Theorem 2.4] *Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form $M = B^+ + C$, where $B^+ = [b_{ij}]$ is the matrix of (4). Then*

$$(1.6) \quad \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \leq \sum_{i=1}^n \frac{n-1}{\min\{\beta_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j},$$

where $\bar{\beta}_i = b_{ii} - \sum_{j=i+1}^n |b_{ij}| l_i(B^+)$ with $l_k(B^+) = \max_{k \leq i \leq n} \left\{ \frac{1}{|b_{ii}|} \sum_{j=k, j \neq i}^n |b_{ij}| \right\}$, and $\prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = 1$ if $i = 1$.

In this paper, we further improve error bounds on the $\text{LCP}(M, q)$ when M belongs to B -matrices. The rest of this paper is organized as follows: In Section 2 we present a new error bound for (1.2), and then prove that this bound are better than those in Theorems 1 and 2. In Section 3, some numerical examples are given to illustrate our theoretical results obtained.

2. A new error bound for the $\text{LCP}(M, q)$ of B -matrices. In this section, an upper bound for (1.2) is provided when M is a B -matrix. Firstly, some definitions, notation and lemmas which will be used later are given as follows.

A matrix $A = [a_{ij}] \in C^{n,n}$ is called a strictly diagonally dominant (SDD) matrix if $|a_{ii}| > \sum_{j \neq i}^n |b_{ij}|$ for all $i = 1, 2, \dots, n$. A matrix $A = [a_{ij}]$ is called a nonsingular M -matrix if its inverse is nonnegative and all its off-diagonal entries are nonpositive [1]. In [6] it was proved that a B -matrix has positive diagonal elements, and a real matrix A is a B -matrix if and only if it can be written in form (1.3) with B^+ being a SDD matrix. Given a matrix $A = [a_{ij}] \in C^{n,n}$, let

$$(2.1) \quad \begin{aligned} w_{ij}(A) &= \frac{|a_{ij}|}{|a_{ii}| - \sum_{\substack{k=j+1, \\ k \neq i}}^n |a_{ik}|}, \quad i \neq j, \\ w_i(A) &= \max_{j \neq i} \{w_{ij}\}, \\ m_{ij}(A) &= \frac{|a_{ij}| + \sum_{\substack{k=j+1, \\ k \neq i}}^n |a_{ik}| w_k}{|a_{ii}|}, \quad i \neq j. \end{aligned}$$

LEMMA 2.1. [19, Theorem 14] *Let $A = [a_{ij}]$ be an $n \times n$ row strictly diagonally dominant M -matrix. Then*

$$\|A^{-1}\|_{\infty} \leq \sum_{i=1}^n \left(\frac{1}{a_{ii} - \sum_{k=i+1}^n |a_{ik}| m_{ki}(A)} \prod_{j=1}^{i-1} \frac{1}{1 - u_j(A) l_j(A)} \right),$$

where $u_i(A) = \frac{1}{|a_{ii}|} \sum_{j=i+1}^n |a_{ij}|$, $l_k(A) = \max_{k \leq i \leq n} \left\{ \frac{1}{|a_{ii}|} \sum_{\substack{j=k, \\ j \neq i}}^n |a_{ij}| \right\}$, $\prod_{j=1}^{i-1} \frac{1}{1 - u_j(A) l_j(A)} = 1$ if $i = 1$, and $m_{ki}(A)$ is defined as in (2.1).

LEMMA 2.2. [15, Lemma 3] *Let $\gamma > 0$ and $\eta \geq 0$. Then for any $x \in [0, 1]$,*

$$\frac{1}{1 - x + \gamma x} \leq \frac{1}{\min\{\gamma, 1\}}$$

and

$$\frac{\eta x}{1 - x + \gamma x} \leq \frac{\eta}{\gamma}.$$

LEMMA 2.3. [16, Lemma 5] Let $A = [a_{ij}]$ with $a_{ii} > \sum_{j=i+1}^n |a_{ij}|$ for each $i \in N$. Then for any $x_i \in [0, 1]$,

$$\frac{1 - x_i + a_{ii}x_i}{1 - x_i + a_{ii}x_i - \sum_{j=i+1}^n |a_{ij}|x_i} \leq \frac{a_{ii}}{a_{ii} - \sum_{j=i+1}^n |a_{ij}|}.$$

Lemmas 2.2 and 2.3 will be used in the proofs of the following lemma and of Theorem 2.5.

LEMMA 2.4. Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form $M = B^+ + C$, where $B^+ = [b_{ij}]$ is the matrix of (1.4). And let $B_D^+ = I - D + DB^+ = [\tilde{b}_{ij}]$ where $D = \text{diag}(d_i)$ with $0 \leq d_i \leq 1$. Then

$$w_i(B_D^+) \leq \max_{j \neq i} \left\{ \frac{|b_{ij}|}{b_{ii} - \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}|} \right\}$$

and

$$m_{ij}(B_D^+) \leq \tilde{m}_{ij}(B^+),$$

where $w_i(B_D^+)$, $m_{ij}(B_D^+)$ are defined as in (2.1), and

$$\tilde{m}_{ij}(B^+) = \frac{1}{b_{ii}} \left(|b_{ij}| + \sum_{\substack{k=j+1, \\ k \neq i}}^n \left(|b_{ik}| \cdot \max_{h \neq k} \left\{ \frac{|b_{kh}|}{b_{kk} - \sum_{\substack{l=h+1, \\ l \neq k}}^n |b_{kl}|} \right\} \right) \right).$$

Proof. Note that

$$[B_D^+]_{ij} = \tilde{b}_{ij} = \begin{cases} 1 - d_i + d_i b_{ij}, & i = j, \\ d_i b_{ij}, & i \neq j. \end{cases}$$

Since B^+ is SDD , $b_{ii} - \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}| > 0$. Hence, by Lemma 2.2 and (2.1), it follows

that

$$(2.2) \quad \begin{aligned} w_i(B_D^+) &= \max_{j \neq i} \{w_{ij}(B_D^+)\} = \max_{j \neq i} \left\{ \frac{|b_{ij}|d_i}{1 - d_i + b_{ii}d_i - \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}|d_i} \right\} \\ &\leq \max_{j \neq i} \left\{ \frac{|b_{ij}|}{b_{ii} - \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}|} \right\}. \end{aligned}$$

Furthermore, it follows from (2.1), (2.2) and Lemma 2.2 that for each $i \neq j$ ($j < i \leq n$)

$$\begin{aligned} m_{ij}(B_D^+) &= \frac{|b_{ij}| \cdot d_i + \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}| \cdot d_i \cdot w_k(B_D^+)}{1 - d_i + b_{ii} \cdot d_i} \\ &\leq \frac{1}{b_{ii}} \cdot \left(|b_{ij}| + \sum_{\substack{k=j+1, \\ k \neq i}}^n |b_{ik}| \cdot w_k(B_D^+) \right) \\ &\leq \frac{1}{b_{ii}} \left(|b_{ij}| + \sum_{\substack{k=j+1, \\ k \neq i}}^n \left(|b_{ik}| \cdot \max_{h \neq k} \left\{ \frac{|b_{kh}|}{b_{kk} - \sum_{\substack{l=h+1, \\ l \neq k}}^n |b_{kl}|} \right\} \right) \right) \\ &= \tilde{m}_{ij}(B^+). \end{aligned}$$

The proof is completed. $\square$

By Lemmas 2.1, 2.2, 2.3 and 2.4, we give the following bound for (1.2) when M is a B -matrix.

THEOREM 2.5. *Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form $M = B^+ + C$, where $B^+ = [b_{ij}]$ is the matrix of (1.4). Then*

$$(2.3) \quad \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \leq \sum_{i=1}^n \frac{n-1}{\min\{\widehat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j},$$

where $\widehat{\beta}_i = b_{ii} - \sum_{k=i+1}^n |b_{ik}| \cdot \tilde{m}_{ki}(B^+)$ with $\tilde{m}_{ki}(B^+)$ is defined in lemma 2.4, $\bar{\beta}_i$ is

defined in Theorem 1.2, and $\prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = 1$ if $i = 1$.

Proof. Let $M_D = I - D + DM$. Then

$$M_D = I - D + DM = I - D + D(B^+ + C) = B_D^+ + C_D,$$

where $B_D^+ = I - D + DB^+ = [\tilde{b}_{ij}]$ and $C_D = DC$. Similarly to the proof of Theorem 2.2 in [10], we can obtain that B_D^+ is an *SDD* M -matrix with positive diagonal elements and that

$$(2.4) \quad \|M_D^{-1}\|_\infty \leq \|(I + (B_D^+)^{-1}\tilde{C}_D)^{-1}\|_\infty \|(B_D^+)^{-1}\|_\infty \leq (n-1)\|(B_D^+)^{-1}\|_\infty.$$

Next, we give an upper bound for $\|(B_D^+)^{-1}\|_\infty$. By Lemma 2.1, we have

$$\|(B_D^+)^{-1}\|_\infty \leq \sum_{i=1}^n \left(\frac{1}{1 - d_i + b_{ii}d_i - \sum_{k=i+1}^n |b_{ik}| \cdot d_i \cdot m_{ki}(B_D^+)} \prod_{j=1}^{i-1} \frac{1}{1 - u_j(B_D^+)l_j(B_D^+)} \right).$$

where

$$u_j(B_D^+) = \frac{\sum_{k=j+1}^n |b_{jk}|d_j}{1 - d_j + b_{jj}d_j}, \quad l_k(B_D^+) = \max_{k \leq i \leq n} \left\{ \frac{\sum_{\substack{j=k, \\ j \neq i}}^n |b_{ij}|d_i}{1 - d_i + b_{ii}d_i} \right\},$$

and

$$m_{ki}(B_D^+) = \frac{|b_{ki}| \cdot d_k + \sum_{\substack{l=i+1, \\ l \neq k}}^n |b_{kl}| \cdot d_k \cdot w_l(B_D^+)}{1 - d_k + b_{kk} \cdot d_k}$$

$$\text{with } w_l(B_D^+) = \max_{h \neq l} \left\{ \frac{|b_{lh}|d_l}{1 - d_l + b_{ll}d_l - \sum_{\substack{s=h+1, \\ s \neq l}}^n |b_{ls}|d_l} \right\}.$$

By Lemmas 2.2 and 2.4, we can easily get that for each $i \in N$,

$$\begin{aligned} \frac{1}{1 - d_i + b_{ii}d_i - \sum_{k=i+1}^n |b_{ik}| \cdot d_i \cdot m_{ki}(B_D^+)} &\leq \frac{1}{\min \left\{ b_{ii} - \sum_{k=i+1}^n |b_{ik}| \cdot m_{ki}(B_D^+), 1 \right\}} \\ &\leq \frac{1}{\min \left\{ b_{ii} - \sum_{k=i+1}^n |b_{ik}| \cdot \tilde{m}_{ki}(B^+), 1 \right\}} \\ (2.5) \quad &= \frac{1}{\min \left\{ \hat{\beta}_i, 1 \right\}}, \end{aligned}$$

and that for each $k \in N$,

$$(2.6) \quad l_k(B_D^+) = \max_{k \leq i \leq n} \left\{ \frac{\sum_{\substack{j=k, \\ j \neq i}}^n |b_{ij}| d_i}{1 - d_i + b_{ii} d_i} \right\} \leq \max_{k \leq i \leq n} \left\{ \frac{1}{b_{ii}} \sum_{\substack{j=k, \\ j \neq i}}^n |b_{ij}| \right\} = l_k(B^+) < 1.$$

Furthermore, according to Lemma 2.3 and (2.6), it follows that for each $j \in N$,

$$(2.7) \quad \frac{1}{1 - u_j(B_D^+) l_j(B_D^+)} = \frac{1 - d_j + b_{jj} d_j}{1 - d_j + b_{jj} d_j - \sum_{k=j+1}^n |b_{jk}| \cdot d_j \cdot l_j(B_D^+)} \leq \frac{b_{jj}}{\bar{\beta}_j}.$$

By (2.5) and (2.7), we have

$$(2.8) \quad \|(B_D^+)^{-1}\|_\infty \leq \frac{1}{\min\{\hat{\beta}_1, 1\}} + \sum_{i=2}^n \left(\frac{1}{\min\{\hat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} \right).$$

The conclusion follows from (2.4) and (2.8). $\square$

The comparisons of the bounds in Theorems 1.2 and 2.5 are established as follows.

THEOREM 2.6. *Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form $M = B^+ + C$, where $B^+ = [b_{ij}]$ is the matrix of (1.4). Let $\bar{\beta}_i$ and $\hat{\beta}_i$ be defined in Theorems 1.2 and 2.5, respectively. Then*

$$\sum_{i=1}^n \frac{n-1}{\min\{\hat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} \leq \sum_{i=1}^n \frac{n-1}{\min\{\bar{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j}.$$

Proof. Note that

$$\bar{\beta}_i = b_{ii} - \sum_{j=i+1}^n |b_{ij}| l_i(B^+), \quad \hat{\beta}_i = b_{ii} - \sum_{k=i+1}^n |b_{ik}| \tilde{m}_{ki}(B^+),$$

and B^+ is a SDD matrix, it follows that for each $i \neq j$ ($j < i \leq n$)

$$\begin{aligned} \tilde{m}_{ij}(B^+) &= \frac{1}{b_{ii}} \left(|b_{ij}| + \sum_{\substack{k=j+1, \\ k \neq i}}^n \left(|b_{ik}| \cdot \max_{h \neq k} \left\{ \frac{|b_{kh}|}{b_{kk} - \sum_{\substack{l=h+1, \\ l \neq k}}^n |b_{kl}|} \right\} \right) \right) \\ &< \frac{1}{b_{ii}} \sum_{\substack{k=j, \\ k \neq i}}^n |b_{ik}| \end{aligned}$$

$$\leq \max_{j \leq i \leq n} \left\{ \frac{1}{b_{ii}} \sum_{\substack{k=j, \\ k \neq i}}^n |b_{ik}| \right\} = l_j(B^+).$$

Hence, for each $i \in N$

$$\widehat{\beta}_i = b_{ii} - \sum_{k=i+1}^n |b_{ik}| \tilde{m}_{ki}(B^+) > b_{ii} - \sum_{k=i+1}^n |b_{ik}| l_i(B^+) = \bar{\beta}_i,$$

which implies that

$$\frac{1}{\min\{\widehat{\beta}_i, 1\}} \leq \frac{1}{\min\{\bar{\beta}_i, 1\}}.$$

This completes the proof. $\square$

Remark here that when $\bar{\beta}_i < 1$ for all $i \in N$, then

$$\frac{1}{\min\{\widehat{\beta}_i, 1\}} < \frac{1}{\min\{\bar{\beta}_i, 1\}},$$

which yields that

$$\sum_{i=1}^n \frac{n-1}{\min\{\widehat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} < \sum_{i=1}^n \frac{n-1}{\min\{\bar{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j}.$$

Next it is proved that the bound (2.3) given in Theorem 2.5 can improve the bound (1.5) in Theorem 1.1 (Theorem 2.2 in [10]) in some cases.

THEOREM 2.7. *Let $M = [m_{ij}] \in R^{n,n}$ be a B -matrix with the form $M = B^+ + C$, where $B^+ = [b_{ij}]$ is the matrix of (1.4). Let β , $\bar{\beta}_i$ and $\widehat{\beta}_i$ be defined in Theorems 1.1, 1.2 and 2.5, respectively, and let $\alpha = 1 + \sum_{i=2}^n \prod_{j=1}^{i-1} \frac{b_{ji}}{\beta_j}$ and $\widehat{\beta} = \min_{i \in N} \{\widehat{\beta}_i\}$. If one of the following conditions holds:*

- (i) $\widehat{\beta} > 1$ and $\alpha < \frac{1}{\bar{\beta}}$;
- (ii) $\widehat{\beta} < 1$ and $\alpha\beta < \widehat{\beta}$,

then

$$\sum_{i=1}^n \frac{n-1}{\min\{\widehat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} < \frac{n-1}{\min\{\beta, 1\}}.$$

Proof. When $\widehat{\beta} > 1$ and $\alpha < \frac{1}{\bar{\beta}}$, we can easily get

$$\sum_{i=1}^n \frac{n-1}{\min\{\widehat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} < \frac{n-1}{\min\{\widehat{\beta}, 1\}} \sum_{i=1}^n \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = (n-1)\alpha < \frac{n-1}{\bar{\beta}} \leq \frac{n-1}{\min\{\beta, 1\}}.$$

Similarly, for $\widehat{\beta} < 1$ and $\alpha\beta < \widehat{\beta}$, the conclusion can be proved directly. $\square$

3. Numerical examples. Two examples are given to show that the bound in Theorem 2.5 is sharper than those in Theorems 1.1 and 1.2.

EXAMPLE 3.1. Consider the family of B -matrices in [15]:

$$M_k = \begin{bmatrix} 1.5 & 0.5 & 0.4 & 0.5 \\ -0.1 & 1.7 & 0.7 & 0.6 \\ 0.8 & -0.1\frac{k}{k+1} & 1.8 & 0.7 \\ 0 & 0.7 & 0.8 & 1.8 \end{bmatrix},$$

where $k \geq 1$. Then $M_k = B_k^+ + C_k$, where

$$B^+ = \begin{bmatrix} 1 & 0 & -0.1 & 0 \\ -0.8 & 1 & 0 & -0.1 \\ 0 & -0.1\frac{k}{k+1} - 0.8 & 1 & -0.1 \\ -0.8 & -0.1 & 0 & 1 \end{bmatrix}.$$

By computations, we have $\beta = \frac{1}{10(k+1)}$, $\bar{\beta}_1 = \bar{\beta}_2 = \frac{90k+91}{100k+100}$, $\bar{\beta}_3 = 0.99$, $\bar{\beta}_4 = 1$, $\hat{\beta}_1 = \frac{820k+828}{900k+900}$, $\hat{\beta}_2 = 0.99$, $\hat{\beta}_3 = 1$ and $\hat{\beta}_4 = 1$. Then it is easy to verify that M_k satisfies the condition (ii) of Theorem 2.7. Hence, by Theorem 1.1 (Theorem 2.2 in [10]), we have

$$\max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \leq \frac{4-1}{\min\{\beta, 1\}} = 30(k+1).$$

It is obvious that

$$30(k+1) \longrightarrow +\infty, \text{ when } k \longrightarrow +\infty.$$

By Theorem 1.2, we have that for any $k \geq 1$,

$$\begin{aligned} & \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \\ & \leq 3 \left(\frac{1}{\bar{\beta}_1} + \frac{1}{\bar{\beta}_2} \cdot \frac{1}{\bar{\beta}_1} + \frac{1}{\bar{\beta}_3} \cdot \frac{1}{\bar{\beta}_1\bar{\beta}_2} + \frac{1}{\bar{\beta}_1\bar{\beta}_2\bar{\beta}_3} \right) \\ & = 3 \left(\frac{100k+100}{90k+91} + \frac{(100k+100)^2}{(90k+91)^2} + \frac{2(100k+100)^2}{0.99(90k+91)^2} \right). \end{aligned}$$

By Theorem 2.5, we have that for any $k \geq 1$,

$$\begin{aligned} & \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \\ & \leq 3 \left(\frac{1}{\hat{\beta}_1} + \frac{1}{\hat{\beta}_2} \cdot \frac{1}{\hat{\beta}_1} + \frac{1}{\hat{\beta}_1\hat{\beta}_2} + \frac{1}{\hat{\beta}_1\hat{\beta}_2\hat{\beta}_3} \right) \end{aligned}$$

$$\begin{aligned}
&= 3 \left(\frac{900k + 900}{820k + 828} + \frac{(100k + 100)}{0.99(90k + 91)} + \frac{1.99(100k + 100)^2}{0.99(90k + 91)^2} \right) \\
&< 3 \left(\frac{100k + 100}{90k + 91} + \frac{(100k + 100)^2}{(90k + 91)^2} + \frac{2(100k + 100)^2}{0.99(90k + 91)^2} \right).
\end{aligned}$$

In particular, when $k = 1$,

$$3 \left(\frac{900k + 900}{820k + 828} + \frac{(100k + 100)}{0.99(90k + 91)} + \frac{1.99(100k + 100)^2}{0.99(90k + 91)^2} \right) \approx 13.9878,$$

$$3 \left(\frac{100k + 100}{90k + 91} + \frac{(100k + 100)^2}{(90k + 91)^2} + \frac{2(100k + 100)^2}{0.99(90k + 91)^2} \right) \approx 14.3775,$$

and the bound (1.5) in Theorem 1.1 is

$$\frac{4 - 1}{\min\{\beta, 1\}} = 30(k + 1) = 60.$$

When $k = 2$,

$$3 \left(\frac{900k + 900}{820k + 828} + \frac{(100k + 100)}{0.99(90k + 91)} + \frac{1.99(100k + 100)^2}{0.99(90k + 91)^2} \right) \approx 14.0265,$$

$$3 \left(\frac{100k + 100}{90k + 91} + \frac{(100k + 100)^2}{(90k + 91)^2} + \frac{2(100k + 100)^2}{0.99(90k + 91)^2} \right) \approx 14.4246,$$

and the bound (1.5) in Theorem 1.1 is

$$\frac{4 - 1}{\min\{\beta, 1\}} = 30(k + 1) = 90.$$

EXAMPLE 3.2. Consider the following family of B -matrices:

$$M_k = \begin{bmatrix} \frac{1}{k} & \frac{-a}{k} \\ 0 & \frac{1}{k} \end{bmatrix},$$

where $\frac{\sqrt{5}-1}{2} < a < 1$ and $\frac{2-a^2}{1+a} < k < 1$. Then $M_k = B_k^+ + C$ with C is the null matrix. By simple computations, we can get

$$\beta = \frac{1-a}{k}, \bar{\beta}_1 = \frac{1-a^2}{k}, \bar{\beta}_2 = \frac{1}{k}, \hat{\beta}_1 = \frac{1}{k} \text{ and } \hat{\beta}_2 = \frac{1}{k}.$$

It is not difficult to verify that M_k satisfies the condition (i) of Theorem 2.7. Thus, the bound (1.6) of Theorem 1.2 (Theorem 2.4 in [17]) is

$$\sum_{i=1}^n \frac{n-1}{\min\{\beta_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = \frac{k+1}{1-a^2},$$

which is larger than the bound

$$\frac{1}{\min\{\beta, 1\}} = \frac{k}{1-a}$$

given by (1.5) in Theorem 1.1 (Theorem 2.2 in [10]). However, by Theorem 2.5 we can get that

$$\max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_{\infty} \leq \frac{2-a^2}{1-a^2}.$$

which is smaller than the bound (1.5) in Theorem 1.1, i.e.,

$$\frac{2-a^2}{1-a^2} < \frac{k}{1-a}.$$

In particular, when $a = \frac{4}{5}$ and $k = \frac{8}{9}$, the bounds in Theorems 1.1 and 1.2 are respectively

$$\frac{1}{\min\{\beta, 1\}} = \frac{k}{1-a} = \frac{360}{81}$$

and

$$\sum_{i=1}^n \frac{n-1}{\min\{\beta_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = \frac{k+1}{1-a^2} = \frac{425}{81},$$

while the bound (2.3) in Theorem 2.5 is

$$\sum_{i=1}^n \frac{n-1}{\min\{\hat{\beta}_i, 1\}} \prod_{j=1}^{i-1} \frac{b_{jj}}{\beta_j} = \frac{2-a^2}{1-a^2} = \frac{306}{81}.$$

These two examples show that the bound in Theorem 2.5 is sharper than those in Theorems 1.1 and 1.2.

Acknowledgements. This work is partly supported by National Natural Science Foundations of China (11601473 and 31600299), Young Talent fund of University Association for Science and Technology in Shaanxi, China (20160234), and CAS 'Light of West China' Program.

REFERENCES

- [1] A. Berman, R.J. Plemmons. *Nonnegative Matrix in the Mathematical Sciences*. SIAM Publisher, Philadelphia, 1994.
- [2] T.T. Chen, W. Li, X. Wu, S. Vong. Error bounds for linear complementarity problems of MB -matrices. *Numer. Algor.*, 70(2):341-356, 2015.

- [3] X.J. Chen and S.H. Xiang. Computation of error bounds for P-matrix linear complementarity problems. *Math. Program., Ser. A* 106:513-525, 2006.
- [4] X.J. Chen and S.H. Xiang. Perturbation bounds of P -matrix linear complementarity problems. *SIAM J. Optim.*, 18:1250-1265, 2007.
- [5] R.W. Cottle, J.S. Pang, R.E. Stone. *The Linear Complementarity Problem*. Academic Press, San Diego, 1992.
- [6] J.M. Peña. A class of P -matrices with applications to the localization of the eigenvalues of a real matrix. *SIAM J. Matrix Anal. Appl.*, 22:1027-1037, 2001.
- [7] P.F. Dai. Error bounds for linear complementarity problems of DB -matrices. *Linear Algebra Appl.*, 434:830-840, 2011.
- [8] P.F. Dai, Y.T. Li and C.J. Lu. Error bounds for linear complementarity problems for SB -matrices. *Numer. Algor.*, 61:121-139, 2012.
- [9] P.F. Dai, C.J. Lu and Y.T. Li. New error bounds for the linear complementarity problem with an SB -matrix. *Numer. Algor.*, 64(4):741-757, 2013.
- [10] M. García-Esnaola and J.M. Peña. Error bounds for linear complementarity problems for B -matrices. *Appl. Math. Lett.*, 22:1071-1075, 2009.
- [11] M. García-Esnaola and J.M. Peña. Error bounds for linear complementarity problems involving B^S -matrices. *Appl. Math. Lett.*, 25(10):1379-1383, 2012.
- [12] M. García-Esnaola and J.M. Peña. Error bounds for the linear complementarity problem with a Σ -SDD matrix. *Linear Algebra and its Applications*, 438(3): 1339-1346, 2013.
- [13] M. García-Esnaola and J.M. Peña. B -Nekrasov matrices and error bounds for linear complementarity problems. *Numer. Algor.*, 72(2):435-445, 2016.
- [14] C.Q. Li, P.F. Dai, Y.T. Li. New error bounds for linear complementarity problems of Nekrasov matrices and B -Nekrasov matrices, *Numer. Algor.*, DOI:10.1007/s11075-016-0181-0.
- [15] C.Q. Li, Y.T. Li. Note on error bounds for linear complementarity problems for B -matrices, *Appl. Math. Lett.*, 57:108-113, 2016.
- [16] C.Q. Li, Y.T. Li. Weakly chained diagonally dominant B -matrices and error bounds for linear complementarity problems, *Numer. Algor.*, DOI: 10.1007/s11075-016-0125-8.
- [17] C.Q. Li, M.T. Gan, S.R. Yang. A new error bound for linear complementarity problems for B -matrices, *Electron. J. Linear Al.*, 31:476-484, 2016.
- [18] K.G. Murty. *Linear Complementarity, Linear and Nonlinear Programming*. Heldermann Verlag, Berlin, 1988.
- [19] Z. Yang, B. Zheng, X. Lian. A New Upper Bound for $\|A^{-1}\|_{\infty}$ of a Strictly α -Diagonally Dominant M -Matrix. *Adv. Numer. Anal.*, 2013:980615, 2013.